

Dietary protein supplementation in the elderly for limiting muscle mass loss

Antonio Herbert Lancha Jr.¹ · Rudyard Zanella Jr.¹ ·
Stefan Gleissner Ohara Tanabe¹ · Mireille Andriamihaja² · Francois Blachier²

Received: 22 September 2016 / Accepted: 18 October 2016 / Published online: 2 November 2016
© Springer-Verlag Wien 2016

Abstract Supplementation with whey and other dietary protein, mainly associated with exercise training, has been proposed to be beneficial for the elderly to gain and maintain lean body mass and improve health parameters. The main objective of this review is to examine the evidence provided by the scientific literature indicating benefit from such supplementation and to define the likely best strategy of protein uptake for optimal objectified results in the elderly. Overall, it appears that an intake of approximately 0.4 g protein/kg BW per meal thus representing 1.2–1.6 g protein/kg BW/day may be recommended taking into account potential anabolic resistance. The losses of the skeletal muscle mass contribute to lower the capacity to perform activities in daily living, emphasizing that an optimal protein consumption may represent an important parameter to preserve independence and contribute to health status. However, it is worth noting that

the maximal intake of protein with no adverse effect is not known, and that high levels of protein intake is associated with increased transfer of protein to the colon with potential deleterious effects. Thus, it is important to examine in each individual case the benefit that can be expected from supplementation with whey protein, taking into account the usual protein dietary intake.

Keywords Whey protein · High protein diet · Anabolic resistance · Protein turnover · Sarcopenia · Aging

Abbreviations

LBM	Lean body mass
MPB	Muscle protein breakdown
mTOR	Mammalian target of rapamycin
MPS	Muscle protein synthesis
IGF-1	Insulin growth factor 1
p70S6k	Protein kinase p70S6
WPI	Whey protein
EAA	Essential amino acids
BCAA	Branched-chain amino acids
FSR	Fractional synthesis rate

Introduction

The human body is in a constant process of protein synthesis and degradation, and this metabolic cycle can be influenced by many factors, including physical activity, caloric deficit and senescence. The rate of renewal is about 300 g of protein per day (Lancha and Pereira-Lancha 2012), representing approximately 2% of the total protein content in adult man weighing 70 kg. Around 40% of body protein are contained in skeletal muscle, and their turnover is considered as relatively slow when compared with the

✉ Antonio Herbert Lancha Jr.
lanchajr@usp.br

✉ Francois Blachier
francois.blachier@agroparistech.fr

Rudyard Zanella Jr.
rudyard.zanella@usp.br

Stefan Gleissner Ohara Tanabe
stefan@usp.br

Mireille Andriamihaja
mireille.andriamihaja@agroparistech.fr

¹ Laboratório de Nutrição e Metabolismo, Escola de Educação Física e Esporte da Universidade de São Paulo, EEFE-USP, R. Prof. Mello Moraes, 65, São Paulo, SP CEP 05508-030, Brazil

² UMR Physiologie de la Nutrition et du Comportement Alimentaire, AgroParisTech, INRA, Université Paris-Saclay, 75005 Paris, France

splanchnic area turnover (Wall and van Loon 2013). The protein turnover is characterized by the balance between muscle protein synthesis (MPS) and muscle protein breakdown (MPB). Resistance training combined with adequate protein intake results in a positive protein turnover, with values of MPS being higher than MPB, leading then to muscle accretion (Cermak et al. 2012). In contrast, caloric restriction periods and/or senescence appears to increase the levels of MPB and lower MPS, resulting in a negative protein turnover and consequently muscle wasting (Areta et al. 2014; Barkoukis 2016; Phillips 2009). Over the time, the human muscle mass is dictated by the changes in MPS and MPB (Deutz and Wolfe 2013; Phillips 2004; Rennie et al. 2004).

To promote a positive protein turnover, it is necessary to take into account the quality of the protein source. Protein derived from dairy foods are considered as among the best in terms of the quality of protein sources (Hartman et al. 2007; Wilkinson et al. 2007). Regarding this latter aspect, whey protein is a fraction (~20%) of the milk protein, which is characterized by a high biological value notably because of its high content in essential amino acids, and particularly high leucine concentration (Pennings et al. 2011a). Leucine has been shown, at least in some cells and animal models, to be able to stimulate protein synthesis through the signaling pathway of the mammalian target of rapamycin complex 1 (mTORC1) (Churchward-Venne et al. 2014a; Dickinson and Rasmussen 2011; Farnfield et al. 2012; Layman et al. 2015). This signaling pathway is responsible for promoting changes in transcription of genes involved in protein synthesis, hence resulting in muscle remodeling (Dickinson and Rasmussen 2011; Farnfield et al. 2012; Fernandes et al. 2008). The mTORC1 is regulated by several cell signaling processes, which can be influenced by many external factors including amino acid availability and physical activity (mainly resistance training) (Churchward-Venne et al. 2014a; Daly et al. 2014; Dickinson et al. 2011; Dickinson and Rasmussen 2011; Kimball and Jefferson 2006; Farnfield et al. 2012). Interestingly, it appears that leucine can interact and stimulate mTORC1 in a different manner than resistance training (Duan et al. 2015; Farnfield et al. 2012; Wolfson et al. 2016; Saxton et al. 2016). Thus, an interesting strategy may be to combine these two stimuli, namely resistance training and leucine availability, to promote protein synthesis through dedicated signaling pathways. This increased stimulation of MPS may result in a positive protein turnover, which can contribute to the maintenance of skeletal muscle mass and possibly for muscle hypertrophy (Churchward-Venne et al. 2014a; Farnfield et al. 2012). Therefore, it is of paramount importance to understand the mechanisms involved in the stimulation of the mTORC1 signaling pathway, since this

pathway is partly responsible for the muscle anabolism (McGlory and Phillips 2015).

The present review aimed at presenting current evidence suggesting that whey and other dietary protein supplementation may exert some beneficial effects in different subpopulations, contributing to the maintenance of skeletal muscle mass in the elderly. Since the upper level of protein consumption with no adverse effect is not known, we also discuss whether dietary protein, when consumed in excessive amounts, may cause some deleterious effects in the large intestine and kidney. To this end, a literature review was conducted using the scientific databases PubMed and Scholar Google.

Dietary protein characteristics

Given the well-known important role of amino acids not only as substrates for protein synthesis, but also as involved in various physiological and metabolic functions, the choice among the different sources of alimentary protein, with different amino acid content and digestibility, represents an important aspect to choose optimal dietary protein for effective effect on skeletal muscle physiology. Regarding this latter point, the whey protein can theoretically guarantee the supply necessary for the stimulation of MPS (Burd et al. 2012; Devries and Phillips 2015; Farnfield et al. 2012; Tang and Phillips 2009). Compared with soy protein, casein and collagen, whey protein has higher relative amount of EAA and leucine. Furthermore, amino acids derived from soy protein for instance appears less bioavailable than amino acids derived from casein and whey protein (Devries and Phillips 2015). In addition, whey protein can release biologically active peptides, which facilitate intestinal absorption (Meisel and Bockelmann 1999). When comparing whey protein with soy protein or whole milk, there is a higher rate of amino acid and oligopeptide absorption in the intestinal tract from the whey protein. Furthermore, whey protein consumption provides a higher stimulation of protein synthesis after resistance exercise when compared to other protein sources (soy and casein) (Poortmans et al. 2012). Lastly, it has been proposed that whey protein may contribute to immune function (Witard et al. 2014).

Recent studies indicate that the availability of intracellular amino acids, particularly leucine, can represent a key regulator of protein synthesis in skeletal muscle after amino acid intake. Previously, it was believed that the pool of extracellular amino acids was the main responsible for this regulation. More studies are required to clarify this issue, but it appears that both intracellular and extracellular amino acids are important for the regulation of protein synthesis (Dickinson and Rasmussen 2011).

Therefore, the increase in the postprandial situation of MPS seems to be modulated by different factors, such as the source of protein ingested (Baer et al. 2011; Phillips et al. 2009; Reitelseder et al. 2011; Walrand et al. 2016), digestibility, postprandial amino acid availability and hyperaminoacidemia/hyperleucinemia (Mitchell et al. 2015a; Smith et al. 1998; Walrand et al. 2016; West et al. 2011; Paddon-Jones et al. 2006; Tang and Phillips 2009; Tipton et al. 1999; Volek et al. 2013; Volpi et al. 2003).

High protein diets and sarcopenia

The quantity and quality of dietary protein have been shown to impact human health parameters in numerous studies (Wu 2016; Blachier et al. 2010). Among the different metabolic and physiological parameters examined, the impact of alimentary protein on skeletal muscle mass loss has been examined (Murphy et al. 2016). The loss of skeletal muscle mass that starts around the 4th decade of life, named sarcopenia, is a process which occurs at a rate of approximately 0.8% per year, accompanied in some cases by a subsequent increase in adipose tissue. To make a long story short, sarcopenia is a multifactorial and complex process resulting from aging-induced changes in the human organism, including altered muscle fiber type distribution and negative net protein balance (Cruz-Jentoft et al. 2010; Fielding et al. 2011; Gumucio and Mendias 2013; Paddon-Jones and Rasmussen 2009; Riddle et al. 2016; Rosenberg 1997). After the sixth decade of life, the loss of skeletal muscle mass may increase to a rate of approximately 15% per decade. Such muscle loss can be worsened by immobility/disuse/bed rest, changes in endocrine function, chronic diseases, inflammation, insulin resistance and nutritional deficiencies (Coker and Wolfe 2012; Fielding et al. 2011; Malafarina et al. 2013). This progressive loss of muscle mass and strength can impair the quality of life, since activities of daily living become more difficult to be performed (da Silva et al. 2014; Velazquez et al. 2013) and the risk of falls and fractures increase (Landi et al. 2012; Scott et al. 2014).

Furthermore, the skeletal muscle is the largest site of lipid oxidation and consumes a large amount of glucose (Argilés et al. 2016; DeFronzo et al. 1992). Then, losses in this tissue result in lower rates of lipid oxidation and an increase in the blood glucose levels, which may contribute to diabetes type II, obesity, and metabolic syndrome (Bauer et al. 2013; Breen and Phillips 2011; Han et al. 2010; Jurca et al. 2005; Ruiz et al. 2008; Wolfe 2006). The skeletal muscle, concomitantly with the liver, is the most important tissue responsible for thermogenesis (Johnstone et al. 2005). Therefore, skeletal muscle plays a key role in the maintenance of metabolic regulation (Argilés et al. 2016; Wolfe 2006).

As a natural geriatric condition, it is important to seek strategies to counteract this process, given the advancing age of the global population, as predictions suggest that about 200 million people will be affected by sarcopenia in 2050 (Janssen 2011). Sarcopenia can be influenced by lifestyle factors, such as reduced physical activity levels (Nair 2005) and suboptimal protein intake (Volpi et al. 2013). It is well established that consuming protein meals containing EAAs and physical activity—mainly resistance exercise—can stimulate MPS (Burd et al. 2010a, b, 2012; Churchward-Venne et al. 2014a; Volek et al. 2013), therefore, serving as key regulators to maintain muscle mass. However, current evidence suggests that, unfortunately, older adults are less sensitive to the stimulatory effects of the nutrient ingestion (Cuthbertson et al. 2005; Pennings et al. 2012) and physical activity (Durham et al. 2010; Fry et al. 2011; Kumar et al. 2009) on MPS compared with younger population.

A possible cause of this less pronounced MPS after nutrient ingestion by the elderly is due to a decreased capacity of digestion and absorption of protein and amino acids (Boirie et al. 1997), concomitantly with a greater amino acids retention by splanchnic area after intestinal absorption (Boirie et al. 1997; Volpi et al. 1999). This may result in less quantities of amino acids available for MPS stimulation. However, there are discrepancies in studies aiming at determining whether the postabsorptive MPS is reduced with aging. Indeed, although several studies have shown lower levels of MPS in the postabsorptive period in the elderly (Welle et al. 1993; Yarasheski et al. 1993); others have shown no significant difference between young and elderly (Cuthbertson et al. 2005; Katsanos et al. 2006). Two recently published cohort studies (Markofskia et al. 2015; Wall et al. 2015) performed over 10 and 5-year periods, respectively, analyzed the basal MPS of over 300 healthy, non-obese young and older adults and showed no difference in the postabsorptive rates of MPS between young and old individuals. These results give an important information indicating that the basal rates of MPS may not be altered with aging to any significant extent.

However, Markofskia et al. (2015) observed a higher phosphorylation of mTOR and its downstream S6K1 in the elderly despite no changes in MPS. This situation was proposed to be associated with an increased mTORC2 signaling and insulin resistance, together with a deficit in the mTOR phosphorylation after the provision of an anabolic stimuli, such as exercise, insulin and nutrition. In agreement with this hypothesis, the decreased sensitivity of MPS may be related to the lower response to insulin (Rasmussen et al. 2006), which may affect both MPS and MPB. Insulin is released to a higher extent by increased amino acids concentrations (Blachier et al. 1989a, b), although, interestingly, old people appears to be less sensitive to this stimuli

(Biolo et al. 1999; Borsheim et al. 2004; Chow et al. 2006; Fujita et al. 2006; Wilkes et al. 2009). This may lead, therefore, to suppressive effects of hyperinsulinemia in the MPS and MPB (Churchward-Venne et al. 2014b; Fujita et al. 2009). In support of this proposition, a possible cause of the impaired MPS in some elderly, as presented by Wall et al. (2015), could be related to the lower insulin sensitivity; this parameter being reflected by higher glycated hemoglobin, fasting insulin, and Homeostatic Model Assessment of Insulin Resistance index.

Moreover, there is also an apparent lower sensitivity of the aging muscle to the stimulation of MPS by hyperaminoacidemia/hyperleucinemia after protein ingestion (Katsanos et al. 2006; Pennings et al. 2012). This may be due to signaling defects, resulting in reduced phosphorylation of mTORC1 and its substrate p70S6k; as a possible consequence of differences in the expression and function of specific amino acids transporters that can be altered with aging (Cuthbertson et al. 2005; Dickinson et al. 2013; Francaux et al. 2016; Guillet et al. 2004).

Increased oxidative stress (Dardevet et al. 2012) and low grade inflammation (Breen and Phillips 2011; Dardevet et al. 2012; Haran et al. 2012; Ticinesi et al. 2016) may also contribute to reduce the ability of some elderly to display a normal stimulation of MPS. Thereby, all these complex mechanisms may be partly responsible for the reduction of the sensitivity of the aging muscle to the stimulatory effects of protein and amino acids ingestion, leading to an impaired capacity to promote muscle protein synthesis. This condition features the so-called “anabolic resistance” of the senescent skeletal muscle (Burd et al. 2013; Rennie 2009), which is responsible for downregulation of MPS which then may be at lower levels than MPB, resulting in a negative protein net balance. Such a negative net balance may lead, over the time, to a loss of skeletal muscle mass. However, all this factors seems to play a secondary role as a cause of anabolic resistance, since the reduced level of habitual physical activity by the older population is likely to be the main responsible for the aggravation observed in this aging condition (Burd et al. 2013; Breen et al. 2013).

The proposed anabolic resistance observed in some elderly is in relationship with the hypothesis of an “anabolic threshold” or “leucine threshold” regulating the MPS, which has to be reached to initiate the MPS (Dardevet et al. 2012; Norton et al. 2009; Rieu et al. 2006). Therefore, it seems that the older population requires greater concentrations of leucine in the blood circulation to achieve this “threshold” for the same MPS than the young individuals (Bauer et al. 2013; Breen and Phillips 2011; Daly et al. 2014; Dardevet et al. 2012; Katsanos et al. 2006). The study performed by Moore et al. (2009) showed that only ~1 g of leucine (about 10 g of whey protein) is necessary to stimulate MPS above basal levels in young individual

muscles at rest, while Yang et al. (2012a) concluded that a minimum amount of ~2 g of leucine (about 20 g of whey protein) are required to rise MPS above basal rates in the elderly at rest.

This issue is of paramount importance as older people tend to consume less dietary protein daily, partly due to reduced energy needs (Fulgoni 2008; Volpi et al. 2013). This remains a serious problem since about one-third of adults who are above the age of 50 consume less than the current recommended dietary allowance (RDA), which is 0.8 g/kgBW/day, whereas approximately 10% of older women do not even reach 0.66 g/kg/BW of protein per day (Houston et al. 2008; Wolfe and Miller 2008; Wolfe et al. 2008). In addition, some elderly ingest small portions of protein in each meal, a fact that contributes to the loss of skeletal muscle mass. Indeed, small quantities of protein seem to have low effects on the stimulation of MPS, even if the quality of protein is high (Katsanos et al. 2005). The situation is further complicated by the fact that older people tend to start the day with small portions of protein (~15 g) and end their day with a meal characterized by high protein content in the evening (~50 g), resulting in a large period of imbalance between MPS and MPB during the day (Berner et al. 2013). This imbalanced protein intake may contribute to decrease the rates of MPS, as the elderly appears to require greater amounts of protein at each meal to raise MPS above basal levels (Moore et al. 2015; Yang et al. 2012a). Then this routine appears to be not ideal for the maintenance of muscle mass with aging, since regular consumption of meals containing protein of high quality (≥ 0.4 g protein/kg BW) throughout the day may likely be more effective to stimulate MPS (Moore et al. 2015; Murphy et al. 2015; Paddon-Jones and Rasmussen 2009).

This latter hypothesis was recently tested by Mamerow et al. (2014) which performed a 7 days crossover study using isonitrogenous diet to measure the changes in MPS over 24 h with an even or irregular protein distribution in healthy young and middle age adults. The even distribution consisted in approximately 30 g of protein in breakfast, lunch and dinner. Over 24-h period, MPS was 25% higher in the even distribution than in the irregular distribution. The strategy of regular protein consumption appears interesting since the rise of MPS by protein intake is transient, reaching the peak within approximately 2 h and returning to basal levels in about 3–4 h after ingestion (Atherton and Smith 2012; Bohe et al. 2001; Dickinson and Rasmussen 2011; Phillips 2014). Furthermore, old people seem to have higher protein needs than young individuals, with values around 1.2–1.6 g protein/kg BW/day (Bauer et al. 2013; Calvani et al. 2013; Wolfe and Miller 2008, 2008). However, the data obtained in this area are rather heterogeneous. Indeed, in a recent publication by Kim et al. (2015), no effect of the protein distribution pattern on the MPS

responses was recorded during an entire day. These findings contrast with the results of Mamerow et al. (2014), maybe because the population studied was different (elderly vs. young/middle age adults) in the two studies. An ingestion of 0.4 g/kg/BW of high-quality protein appears to be the minimum amount required to stimulate muscle anabolic response (Moore et al. 2015).

When ingested in the context of a mixed meal, it is possible that a greater amount of high-quality protein is required to achieve the maximal MPS, as this condition is associated with less amino acids bioavailability (Burke et al. 2012). Nevertheless, it still can be recommended to fractionate the protein intake during the day, since older people frequently feature a decreased level of appetite (Landi et al. 2016), a condition that may hamper the consumption of a meal with very high protein content, as seen in the study by Kim et al. (2015). It is worth noting that the main difference between the higher vs. lower protein ingestion groups was the increased rates of MPS in the higher protein ingestion group, a result in accordance with other findings (Bauer et al. 2013; Calvani et al. 2013; Wolfe and Miller 2008, 2008) and which suggests that older individual may benefit from higher consumption of protein during the all day.

A recent work (Loenneke et al. 2016) found a positive association between protein quantity per meal and leg lean mass and strength. In fact, consuming frequent meals containing 30–45 g of protein is associated with greater leg lean mass and knee extensor muscle strength. Such nutritional strategy was also associated with higher LBM and appendicular LBM in both older men and women at baseline and after a 2-years follow-up period (Farsijani et al. 2016). These findings reinforce those provided by Loenneke et al. (2016). However, further long-term longitudinal research is required to determine an optimal protein intake quantity and distribution throughout the day to help to preserve skeletal muscle mass and function in older individuals.

The difficulty to reach the optimal value of protein intake during the day by the elderly may lead to a protein deficit, which chronically may induce skeletal muscle wasting. To counteract this situation, high protein diets combined with high-quality protein supplement may help for the preservation of skeletal muscle mass (Cermak et al. 2012; Deer and Volpi 2015; Malafarina et al. 2013; Wolfe 2012). To support this proposition, some studies (Geirsdotir et al. 2013; Houston et al. 2008) have found a positive correlation between consumption of protein and maintenance of skeletal muscle mass in older adults, since higher amounts of protein intake was correlated with an increased skeletal muscle mass retention over the time; suggesting a possible protective effect of protein against muscle wasting when amounts of protein near or even above the recommended values are ingested.

Consumption of whey protein instead of protein meals has been proposed as a possible strategy to reach the daily requirements of protein able to stimulate MPS in elderly. A study performed by Pennings et al. (2012) compared the rates of protein accretion after the ingestion of three different doses of whey protein (10, 20, 35 g) and showed positive results for the 20 and 35 g-consuming groups while the MPS displayed no significant difference after the ingestion of the 10 g protein dose. This result is in agreement with the findings from Wall et al. (2015) which suggest that higher doses of protein with high biological value at rest are necessary to optimally stimulate MPS in older men. It may explain why Arnal et al. (1999) and Bouillanne et al. (2013) did not find positive results under their experimental design. Indeed, Bouillanne et al. (2013) fail to demonstrate any benefit from 6 weeks dietary intervention in hospitalized older adults, with most meals containing less than 20 g protein. Similar results were obtained in the study by Arnal et al. (1999). Thus, it appears that 20 g protein is the minimal amount required to promote MPS in elderly at rest (Yang et al. 2012a).

Positive results have been obtained by Tieland et al. (2012b) after the intake of the supplement in breakfast and lunch of frail elderly. In this latter study, protein intake increased to more than 25 g of protein in each main meal (≥ 0.34 g/kg/BW per meal), allowing improvements of both strength and physical performance despite no increase in skeletal muscle mass after 24 weeks.

In a similar manner, an optimized offer of high-quality protein in the breakfast and lunch (all daily meals containing ≥ 0.4 g/kg/BW) by protein supplementation promotes a significant increase in appendicular lean tissue mass in healthy older individuals (Norton et al. 2016). In this latter and other studies, the average compliance to the treatment was very high ($\geq 92\%$) (Norton et al. 2016; Tieland et al. 2012b). In addition, since the protein-induced suppression of energy intake may be blunted in the elderly (Giezenaar et al. 2015; Norton et al. 2016), this may explain why the maintenance of energy balance and consequently of a positive skeletal muscle protein balance were observed (Carbone et al. 2012; Hector et al. 2015; Pasiakos et al. 2015).

Physical activity may also play an import role for the maintenance of skeletal muscle mass and function. Resistance training combined with milk protein supplementation given in the breakfast and lunch of frail elderly gave promising results since this combined stimuli increases by 1.3 kg lean body mass with no differences in the placebo group (Tieland et al. 2012a). However, similar to others findings (Norton et al. 2016; Tieland et al. 2012b), and as pointed out above, there was an increase in the total daily protein intake from 1.0 g/kg/BW up to 1.3 g/kg/BW, rendering impossible to attribute these results only to the protein distribution pattern.

Yang and colleagues (2012a), by combining resistance exercise in older men with post-exercise whey protein supplementation, found that different doses of protein, i.e., 10 g (0.13 g/kg BW), 20 g (0.25 g/kg BW) and 40 g (0.49 g/kg BW) gave positive results. They found that, as observed by Pennings et al. (2012) who demonstrated better results in MPS with higher doses of protein ingestion at rest; 40 g of whey protein intake given after resistance training was able to stimulate myofibrillar protein synthesis by 91% above the placebo group; while the group that consumes 20 g of whey displayed only 44% increase above the value obtained in volunteers without supplementation. In agreement, D'Souza et al. (2014) found a linear relationship between whey protein dose and muscle p70S6k phosphorylation, wherein 40 g of protein intake results in greater levels of p70S6k phosphorylation than lower protein doses. Phosphorylation of p70S6k is often correlated to increases in MPS rates and possible muscle hypertrophy (Baar and Esser 1999; Mitchell et al. 2013; Terzis et al. 2008). Although, other mechanisms are likely to be also involved in the skeletal muscle hypertrophy. In fact, in some cases there is a lack of correlation between acute changes in phosphorylation of p70S6k and long-term skeletal muscle hypertrophy (Mitchell et al. 2012, 2014). These findings provide an important information suggesting that older people have higher need for high-quality protein to optimize the stimuli of muscle protein synthesis at rest and after resistance exercise.

In agreement with these results, a recently published randomized, double-blind, placebo-controlled supplementation trial (Rondanelli et al. 2016) that combined whey protein, essential amino acids, and vitamin D with regular physical activity in 130 sarcopenic elderly people (53 men and 77 women; mean age 80.3 years) showed positive results of the dietary intervention. After 12 weeks of intervention with physical activity 5 times/week and ingestion of 32 g of a supplement containing 22 g of whey protein (~4 g of leucine) and 2.5 g of vitamin D one time per day, 68% of the sarcopenic elderly became non-sarcopenic with a gain of ~1.7 kg in fat free mass. These results were concomitant with others improvements in health parameters, such as reduced inflammatory state, increased performance on activities of daily living, and enhanced IGF-1 concentrations. These results suggest that physical activity is important, but alone is not sufficient, to achieve significant results since placebo group did not present all these improvements. However, the physical activity was non intensive in this latter study, suggesting that it could have been a limiting factor for the increase in fat free mass. Furthermore, it is important to notice that despite the relatively low dose of whey protein used (22 g), the high content of these protein in leucine and the vitamin D supplementation might explain the positive and ample results of the study.

Current evidence suggests that performing exercise before protein intake allows greater use of dietary protein-derived amino acids for de novo muscle protein synthesis in elderly men (Pennings et al. 2011b). Thus, based on these data, physical activity positively impacts the ability of skeletal muscle to retain dietary amino acids, in both young and old individuals, and improves the anabolic response of a meal containing protein (Devries et al. 2015; Timmerman et al. 2012; Walker et al. 2011). A bout of moderate intensity aerobic exercise seems to be efficient to sensitize the skeletal muscle and enhances the anabolic effects of protein feeding (Timmerman et al. 2012). Although the enhanced sensitivity of MPS after an intense bout of resistance exercise persists up to 24 h (Damas et al. 2015), the greatest exercise-mediated increases in MPS occurs immediately after the exercise bout (Churchward-Venne et al. 2012). Thus, providing frequent daily stimulus such as walking to enhance MPS may be valuable for the elderly to maintain the maximal skeletal muscle response to protein-induced increases in MPS during the day.

Due to anabolic resistance, as defined above, an important issue that thus appears determinant for the stimulation of skeletal muscle protein synthesis is the protein source. Indeed, there are different types of dietary proteins with different characteristics, such as the digestibility, amino acids composition, and rates and kinetics of absorption of these AAs. Such parameters would likely lead to distinct MPS responses. Data provided by Yang et al. (2012b) compared the MPS responses after intake of different protein sources in the elderly. These authors assessed the effects of whey protein and soy protein ingestion under rest and post-exercise conditions. They compared three different doses of protein supplementation, that are 0, 20 and 40 g of whey protein or soy protein on the rates of leucine oxidation and fractional synthesis rates (FSR). The results pointed to higher increases in FSR in both rested and after resistance exercise using whey in comparison with soy protein, despite lower rates of leucine oxidation when small doses of protein were ingested. Similar results were found by Mitchell et al. (2015a), which showed a less prolonged p70S6k phosphorylation after soy protein ingestion (~2 h) compared to whey protein (~4 h) after a session of resistance exercise.

Whey protein also seems to be more effective to increase the rates of MPS when compared with casein. A study performed by Burd et al. (2012) revealed that the ingestion of whey protein elicited greater rates of MPS against its casein counterpart in older adults. It is important to notice that this latter study also tested the subjects at rest and after resistance training, and found, in both conditions, whey to give better results than casein. This result is probably related to a greater hyperaminoacidemia or hyperleucinemia after whey ingestion, due to the kinetic of digestion

of each protein. Findings from Pennings et al. (2011a, b) is in accordance with these results, as greater circulating concentrations of leucine were measured after whey ingestion compared with casein ingestion. Moreover, high leucine plasma concentrations combined with peak leucinemia resulting from whey supplementation displayed a strong relationship with MPS (Pennings et al. 2011a, b). Walrand et al. (2016) recently compared in the elderly the rates of synthesis in individual muscle protein fraction—sarco-plasmic, mitochondrial, actin and myosin—after the intake of casein and a fast-digestive milk protein with different amounts of leucine. The consumption of both 15 and 30 g of the fast-digestive milk protein was able to increase mitochondrial protein FSR, while no difference was elicited in the postprandial period after casein ingestion, suggesting that each protein source has an unique effect at specific muscle protein fractions.

However, although myosin protein FSR was stimulated after both protein sources, myosin FSR was enhanced with 30 g but not with 15 g of casein. Sarcoplasmic muscle protein and actin FSRs were not significantly increased in the postprandial state compared to the postabsorptive state. It is worth to note that the availability of plasma leucine was higher for both 15 and 30 g of fast-digestive milk protein, despite 30 g of casein contain larger amounts of leucine than 15 g of rapidly digested milk protein.

These results point to a better response of MPS after whey ingestion despite the fact that casein (only when hydrolyzed) and soy protein also display fast digestibility, high bioavailability, and high EAAs content. Whey protein consumption results in a higher concentrations of leucine and promotes an overall rapid increase of amino acid concentrations in plasma. Such increase apparently is the “trigger” to start the cell signaling process allowing protein synthesis (Katsanos et al. 2006; Layman et al. 2015; Volpi et al. 2003; West et al. 2011). Concomitantly, the role of each source of protein in the human metabolism may contribute to the differences in the MPS stimulation after the ingestion of a given mixture of nutrients. Indeed, for instance, amino acid derived from whey protein in milk or from soy protein can be used differently both in


the splanchnic and peripheral tissues. Amino acids derived from soy protein seems to support greater splanchnic protein synthesis and to be converted to urea to a greater extent than milk proteins. This may be due to the high content of BCAAs in whey protein, since it contributes to less splanchnic extraction (Bos et al. 2003; Fouillet et al. 2002; Luiking et al. 2005). These different metabolic fates may explain, in part at least, the superiority of milk protein to rise MPS compared to soy protein (Hartman et al. 2007; Wilkinson et al. 2007).

Without consideration regarding the different dietary sources, high quantities of protein may be efficient to stimulate MPS above basal rates when combined with resistance training in elderly. Some studies have demonstrated that 40 g of beef meat, whey and soy protein, were able to raise MPS around 80–100% after resistance training. However, low doses (~10 g) were unable to increase the protein accretion (Robinson et al. 2013; Yang et al. 2012a, b).

However, in contrast to the findings above, two recently published data in both young individuals (Mitchell et al. 2015b) and in the elderly (Mitchell et al. 2015c) suggest that a sustained EAA delivery profile (pulse feeding) may be as good as a rapid rise of aminoacidemia (bolus feeding) to stimulate MPS at rest. More studies applying this latter strategy are needed to explore potential clinical benefits of such a strategy for the elderly.

Taken together, these findings show that most elderly people do not reach the necessary amounts of protein to counteract the anabolic resistance and to attenuate the subsequent sarcopenia. A strategy that seems to be useful to help old people to ingest their protein daily requirements is through protein supplementation, mainly in the form of whey protein due to their relatively high EAAs content (notably with high amounts of leucine), fast and high digestibility and good bioavailability. All these characteristics are propitious to promote a rapid and robust rise in leucinemia, likely allowing high rates of MPS. Thus, providing three or more high-quality, leucine-rich protein meals during the day combined with regular physical activity may represent a good strategy to attenuate aging-related skeletal muscle mass losses (Fig. 1).

Fig. 1 Schematic view showing how the key variables of protein ingestion and physical activity can influence the skeletal muscle mass in the elderly. *MPS* muscle protein synthesis, *MPB* muscle protein breakdown



High protein diet and renal impairment

The high protein intake leads to oxidation of the amino acids in excess and possible metabolism in triglycerides and glucose. Excessive protein intake results in an elevated nitrogen rates in the body, which leads to a greater glomerular filtration rate, and increases the production of urea (Marckmann et al. 2015; Poortmans et al. 2012). There was no effect of a high protein intake on renal function, such as glomerular filtration, excretion of albumin and calcium metabolism in trained athletes consuming less than 2.8 g/kgBW per day of protein (Poortmans and Dellalieux 2000). In another study, the renal function was examined by the measurement of the glomerular filtration rate, after 12 weeks of intervention with resistance training and protein supplementation in elderly. The results showed no impairment of glomerular filtration rate with a protein consumption of approximately 1.0 g/kgBW (Ramel et al. 2013).

The current recommendation suggests a moderate/restricted protein consumption, mainly in the elderly, only in the presence of a moderate/severe kidney disease with no dialysis (Bauer et al. 2013).

High-protein diet and colon health

An increase in the amount of dietary protein intake increases the quantity of unabsorbed proteins, peptides and amino acids reaching the large intestine through the ileocecal junction (Gibson et al. 1976). In the small and large intestines, bacteria in the luminal contents can metabolize proteins and amino acids (Dai et al. 2012; Blachier et al. 2010; Andriamihaja et al. 2013); and metabolomics analysis has revealed different profiles in the plasma and urine of rodents receiving a high-protein diet as compared to diet containing a moderate amount of proteins (Mu et al. 2015). In the large intestine, undigested and partially digested dietary protein and peptides from dietary and endogenous origin are degraded by endogenous and bacterial proteases/peptidases in peptides and amino acids; these latter being not absorbed by the colonic epithelium but serving as precursors for numerous bacterial metabolites including ammonia, hydrogen sulfide, amines, short-chain and branched-chain fatty acids, indoles, phenols, ethanol and organic acids, etc. (Liu et al. 2014; Mouillé et al. 2004; Davila et al. 2013). Several among these metabolites (H_2S , NH_4^+ , *p*-cresol) have been shown at excessive concentrations to affect colonic epithelial energy metabolism (Beamont et al. 2016; Andriamihaja et al. 2010, 2015). *p*-Cresol which is produced by the microbiota from *L*-tyrosine shows genotoxic effect on colonocytes (Andriamihaja et al. 2015). In addition, hydrogen sulfide has been recently shown to drive mucin denaturation, and to presumably reduce mucus barrier function

in the colon (Ijssennagger et al. 2015). High-protein diet given for 2 weeks has been shown to modify colonic epithelial cell morphology (Andriamihaja et al. 2010) and to affect the distribution of mucous cells in the colonic epithelium (Lan et al. 2015), maybe due to alteration in the process of colonic epithelial cell proliferation and differentiation. These results were all obtained in animal models. In human studies, it has been shown that high protein weight loss diet promotes in the colonic lumen a bacterial metabolite profile that is likely detrimental for colonic health (Russell et al. 2011); suggesting that long-term adherence to such diet may increase risk of colonic diseases. Since there is no upper limit for protein intake with no adverse effects, some caution is advised regarding long-term consumption of high protein diet particularly regarding the maintenance of the colonic health.

Conclusions

From the available data, it appears necessary in the elderly population to examine each case individually to determine the actual relevance for protein dietary supplementation. Furthermore, one must take into account the origin of the substances that will be used for supplementation, as well as the safety for use and evidence of its effectiveness.

A key point regarding the effectiveness of protein supplementation is the combination of training, food intake and supplementation. Considering that muscle hypertrophy is only due to protein supplementation has simply no scientific support. Several studies showed some positive results on the association of physical exercise together with supplementation of whey protein, especially for audiences like the elderly who may experience protein deficit. It is worth to note that protein sources with high biological value containing essential amino acids (notably leucine), such as whey protein, but of course not exclusively, are most relevant to generate the appropriate stimulus for protein synthesis. The Table 1 presents a list of studies related to the effects of dietary protein supplementation in the elderly.

Regarding the recommended dose, an adequate protein intake from 1.2 up to 1.6 g/kg/BW during the day may be required by the elderly to preserve muscle mass. Fractionate protein intake in daily doses may likely be a good strategy since older individuals may experience diminished appetite. Therefore, an individualized nutritional strategy should always be designed to successfully reach the recommended daily protein needs. However, since the upper non deleterious level of dietary protein remains undefined, and taking into consideration that high protein diet in long-term utilization may impact colon health, some additional studies are required to test in healthy volunteers the impact of high-protein diet on the large intestine epithelium. In that

Table 1 Summary of the quoted studies related to the effects of dietary protein supplementation in aging individuals and in the elderly

Study	Duration	Subjects	Protocol	Outcome
Burd et al. (2012)	Acute	14 healthy aging men	Ingestion of 20 g of casein or whey at rest and after an acute bout of RE	Greater myofibrillar MPS after whey than casein in both conditions
Penning et al. (2012)	Acute	33 healthy aging men	Ingestion of 10, 20 and 35 g of whey protein at rest	Muscle protein accretion 35 g > 20 g > 10 g
Wall et al. (2015)	Acute, retrospective, cross sectional study	35 healthy young men; 40 healthy elderly men	Ingestion of 20 g of casein at rest	16% lower post-prandial FSR in elderly than in young men
Francaux et al. (2016)	Acute	9 healthy young men; 10 healthy aging men	Ingestion of 30 g of whey protein after an unilateral RE	↑ PKB phosphorylation in young than elderly in both rested and after RE ↑ mTOR in both age groups after RE ↓ S6K1 in all conditions in the aging men
Yang et al. (2012b)	Acute	30 healthy elderly men	Ingestion of 0, 20, 40 g of whey protein (WP) or soy protein (SP) at rest and after a bout of RE	Leucine oxidation: SP20 > WP20; SP40 = WP40 MPS at rest: SP0 = SP20 = SP40; WP40 > WP20 > WP0 MPS after RE: SP40 > SP0 = SP20; WP40 > SP40 = WP20
Moore et al. (2015)	Acute	65 healthy young men; 43 healthy aging men	Ingestion of varying amounts (0–40 g) of high-quality dietary protein as a single bolus and normalized to body mass	No differences in basal MPS between the groups MPS reached a plateau after ingestion of 0.40 g/kg/BW and 0.24 g/kg/BW in aging and young men, respectively
D'Souza et al. (2014)	Acute	46 healthy elderly untrained men	Ingestion of 10, 20, 30 or 40 g of whey protein or placebo after a bout of RE	↓ Intramuscular BCAAs levels after RE in placebo group ↑ Intramuscular BCAAs levels after 2 h post RE in the 30 and 40 g whey protein groups, and after 4 h only in 40 g whey protein group
Mitchell et al. (2015c)	Acute	33 healthy aging untrained men	Ingestion of 30 g of whey protein, soy protein, CHO or placebo	Significantly increases in p70S6K phosphorylation up to 4 h only after ingestion of ≥20 g of whey protein. Best results after 30 and 40 g of whey protein Increased p70S6K in whey and soy protein group after 2 h post RE After 4 h post RE p70S6K remained elevated only in whey protein group
Penning et al. (2011a)	Acute	48 aging men	Ingestion of 20 g whey, casein or casein hydrolysate	FSR values: whey > casein hydrolysate > casein
Penning et al. (2011b)	Acute	24 young and 24 aging men	Ingestion of 20 g casein at rest or after a 30 min exercise protocol	Greater use of AAs for MPS after exercise in young and elderly men
Farnfield et al. (2012)	Acute and chronic (12 weeks)	16 healthy young men; 15 healthy aging men	Ingestion of 27 g of whey protein or placebo after an acute bout of RE pre and post 12 week RE program	Similar increases in strength in both groups for each category Pre RE: ↑ mTOR, 4E-BP1 after whey than placebo in both ages ↑↑ p70S6K after whey in older compared with placebo ↑ eIF4G after whey than placebo in older Post RE: slightly blunted mTOR pathway phosphorylation after whey ingestion in aging men

Table 1 continued

Study	Duration	Subjects	Protocol	Outcome
Walrand et al. (2016)	Chronic, 10 days	31 healthy elderly men	Ingestion of 15 g adequate protein (AP) or 30 g high-protein (HP) diets of Casein (CAS) or soluble Milk protein (PRO)	↑ Myosin FSR after both HP groups, but only in PRO AP ↑ Mitochondrial FSR in both PRO quantities
Norton et al. (2016)	Chronic, 24 weeks	60 healthy ageing men and women	Ingestion of 0.165 g/kg body mass of a milk-based protein matrix (PRO) or an isoelectric, non-nitrogenous maltodextrin control (CON) at breakfast and midday meals	Increased daily protein ingestion from 1.2 to 1.6 g/kg/BW in the PRO group Protein intakes in the PRO group increased to ≥ 0.4 g/kg per meal LTM increased by 0.45 kg in PRO group and decreased 0.16 kg in CON group
Tieland et al. (2012b)	Chronic, 24 weeks	65 pre-frail and frail elderly men and women	Ingestion of 15 g of milk protein (PRO) or placebo after breakfast and lunch	Increased daily protein ingestion from 1 to 1.4 g/kg/BW in the PRO group Protein intake in the PRO group increased to more than 25 g per meal Improved physical performance in PRO group compared with placebo group
Tieland et al. (2012a)	Chronic, 24 weeks	62 frail elderly subjects	Ingestion of 15 g of milk protein (PRO) or placebo after breakfast and lunch Participation in a progressive resistance-type exercise training program 2 × per week	Increased daily protein ingestion from 1 g/kg/BW to 1.3 g/kg/BW in the PRO group ↑ LBM in protein group, no changes in placebo group Strength and physical performance improved significantly in both groups
Rondanelli et al. (2016)	Chronic, 12 weeks	130 sarcopenic elderly (53 men; 77 women)	Ingestion of nutritional supplement (22 g whey protein, 4 g leucine, 2.5 g vitamin D) or placebo Both groups participated in a five times/week physical activity program	Fat free mass increased 1.7 kg in the supplemented group; no changes observed in placebo group Significant improvements in relative skeletal muscle mass, strength, inflammatory state and physical function in the supplemented group; no changes observed in placebo group 68% of elderly in the supplemented group became non-sarcopenic

RE resistance exercise, MPS muscle protein synthesis, FSR fractional synthesis rate, PKB protein kinase B, mTOR mammalian target of rapamycin, p70S6K1 or S6K1 ribosomal protein S6 kinase 1, BCAAs branched-chain amino acids, AAs amino acids, 4E-BP1 eukaryotic translation initiation factor 4E-binding protein 1, eIF4G eukaryotic translation initiation factor 4G, LTM lean tissue mass, LBM lean body mass, FFM fat free mass, RMM relative skeletal muscle mass

regard, supplementation with individual amino acids may represent on some occasions an interesting strategy, taking into account the large capacity of the small intestine for their absorption.

Acknowledgements The authors wish to thank the University of Sao Paulo, AgroParisTech, INRA and Université Paris-Saclay for their support.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethics statement The manuscript has not been submitted to other journal for simultaneous consideration. The manuscript, partly or in full, has not been published previously. No data have been fabricated or manipulated (including images) to support our conclusions. No data, text, or theories by others are presented as if they were the author's own (plagiarism).

References

- Andriamihaja M, Davila AM, Eklou-Lawson M et al (2010) Colon luminal content and epithelial cell morphology are markedly modified in rats fed with a high-protein diet. *Am J Physiol Gastrointest Liver Physiol* 299:G1030–G1037
- Andriamihaja M, Guillot A, Svendsen A et al (2013) Comparative efficiency of microbial enzymes preparations versus pancreatin for in vitro alimentary protein digestion. *Amino Acids* 44:563–572
- Andriamihaja M, Lan A, Beaumont M et al (2015) The deleterious metabolic and genotoxic effects of the bacterial metabolite p-cresol on colonic epithelial cells. *Free Radic Biol Med* 85:219–227
- Areta J, Burke L, Camera D et al (2014) Reduced resting skeletal muscle protein synthesis is rescued by resistance exercise and protein ingestion following short-term energy deficit. *Am J Physiol Endocrinol Metab* 306:E989–E997
- Argilés JM, Campos N, Lopez-Pedrosa JM et al (2016) Skeletal muscle regulates metabolism via interorgan crosstalk: roles in health and disease. *J Am Med Dir Assoc* 17(9):789–796
- Arnal MA, Mosoni L, Boirie Y et al (1999) Protein pulse feeding improves protein retention in elderly women. *Am J Clin Nutr* 69:1202–1208
- Atherton PJ, Smith K (2012) Muscle protein synthesis in response to nutrition and exercise. *J Physiol* 590(5):1049–1057
- Baar K, Esser K (1999) Phosphorylation of p70(S6k) correlates with increased skeletal muscle mass following resistance exercise. *Am J Physiol* 276:C120–C127
- Baer DJ, Stote KS, Paul DR et al (2011) Whey protein but not soy protein supplementation alters body weight and composition in free-living overweight and obese adults. *J Nutr* 141(8):1489–1494
- Barkoukis H (2016) Muscle building and maintenance in the elderly: the use of protein. *Curr Nutr Rep*. doi:10.1007/s13668-016-0163-9
- Bauer J et al (2013) Evidenced-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. *J Am Med Dir Assoc* 14:542–559
- Beaumont M, Andriamihaja M, Lan A et al (2016) Detrimental effects for colonocytes of an increased exposure to luminal hydrogen sulfide: the adaptive response. *Free Radic Biol Med* 93:155–164
- Berner L, Becker G, Wise M et al (2013) Characterization of dietary protein among older adults in the United States: amount, animal sources, and meal patterns. *J Acad Nutr Diet* 113:809–815
- Biolo G, Williams BD, Fleming RY et al (1999) Insulin action on muscle protein kinetics and amino acid transport during recovery after resistance exercise. *Diabetes* 48(5):949–957
- Blachier F, Leclercq-Meyer V, Marchand J et al (1989a) Stimulus-secretion coupling of arginine-induced insulin release. Functional response of islets to L-arginine and L-ornithine. *Biochim Biophys Acta* 1013:144–151
- Blachier F, Mourtada A, Sener A et al (1989b) Stimulus-secretion coupling of arginine-induced insulin release. Uptake of metabolized and non-metabolized cationic amino acids by pancreatic islets. *Endocrinology* 124:134–141
- Blachier F, Lancha AH Jr, Boutry C et al (2010) Alimentary proteins, amino acids, and cholesterolemia. *Amino Acids* 38:15–22
- Bohe J, Low JF, Wolfe RR et al (2001) Latency and duration of stimulation of human muscle protein synthesis during continuous infusion of amino acids. *J Physiol* 532:575–579
- Boirie Y, Gachon P, Beaufrere B (1997) Splanchnic and whole-body leucine kinetics in young and elderly men. *Am J Clin Nutr* 65:489–495
- Borsheim E, Cree MG, Tipton KD et al (2004) Effect of carbohydrate intake on net muscle protein synthesis during recovery from resistance exercise. *J Appl Physiol* 96:674–678
- Bos C, Metges CC, Gaudichon C et al (2003) Postprandial kinetics of dietary amino acids are the main determinant of their metabolism after soy or milk protein ingestion in humans. *J Nutr* 133:1308–1315
- Bouillanne O, Curis E, Hamon-Vilcot B et al (2013) Impact of protein pulse feeding on lean mass in malnourished and at-risk hospitalized elderly patients: a randomized controlled trial. *Clin Nutr* 32:186–192
- Breen L, Phillips SM (2011) Skeletal muscle protein metabolism in the elderly: interventions to counteract the ‘anabolic resistance’ of ageing. *Nutr Metab* 8:68
- Breen L, Stokes KA, Churchward-Venne TA et al (2013) Two weeks of reduced activity decreases leg lean mass and induces “anabolic resistance” of myofibrillar protein synthesis in healthy elderly. *J Clin Endocrinol Metab* 98:2604–2612
- Burd NA, Holwerda AM, Selby KC et al (2010a) Resistance exercise volume affects myofibrillar protein synthesis and anabolic signalling molecule phosphorylation in young men. *J Physiol* 588:3119–3130
- Burd NA, West DW, Staples AW et al (2010b) Low-load high volume resistance exercise stimulates muscle protein synthesis more than high-load low volume resistance exercise in young men. *PLoS One* 5:e12033
- Burd NA, Yang Y, Moore DR et al (2012) Greater stimulation of myofibrillar protein synthesis with ingestion of whey protein isolate v. micellar casein at rest and after resistance exercise in elderly men. *Br J Nutr* 108:958–962
- Burd NA, Gorissen SH, van Loon LJC (2013) Anabolic resistance of muscle protein synthesis with aging. *Exerc Sport Sci Rev* 41(3):169–173
- Burke LM, Winter JA, Cameron-Smith D et al (2012) Effect of intake of different dietary protein sources on plasma amino acid profiles at rest and after exercise. *Int J Sport Nutr Exerc Metab* 22:452–462
- Calvani R, Miccheli A, Landi F et al (2013) Current nutritional recommendations and novel dietary strategies to manage sarcopenia. *J Frailty Aging* 2(1):38–53
- Carbone JW, McClung JP, Pasiakos SM (2012) Skeletal muscle responses to negative energy balance: effects of dietary protein. *Adv Nutr* 3:119–126

- Cermak NM, Res PT, Groot LC et al (2012) Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. *Am J Clin Nutr* 96(6):1454–1464
- Chow LS, Albright RC, Bigelow ML et al (2006) Mechanism of insulin's anabolic effect on muscle: measurements of muscle protein synthesis and breakdown using aminoacyl-tRNA and other surrogate measures. *Am J Physiol Endocrinol Metab* 291:E729–E736
- Churchward-Venne TA, Burd NA, Phillips SM (2012) Nutritional regulation of muscle protein synthesis with resistance exercise: strategies to enhance anabolism. *Nutr Metab (Lond)* 9:40
- Churchward-Venne TA, Breen L, Di Donato DM et al (2014a) Leucine supplementation of a low-protein mixed macronutrient beverage enhances myofibrillar protein synthesis in young men: a double-blind, randomized trial. *Am J Clin Nutr* 99(2):276–286
- Churchward-Venne TA, Breen L, Phillips SM (2014b) Alterations in human muscle protein metabolism with aging: protein and exercise as countermeasures to offset sarcopenia. *BioFactors* 40(2):199–205
- Coker RH, Wolfe RR (2012) Bedrest and sarcopenia. *Curr Opin Clin Nutr Metab Care* 15:7–11
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM et al (2010) Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *European Working Group on Sarcopenia in Older People. Age Ageing* 39:412–423
- Cuthbertson D, Smith K, Babraj J et al (2005) Anabolic signaling deficits underlie amino acid resistance of wasting, aging muscle. *FASEB J* 19:422–424
- D'Souza RF, Markworth JF, Figueiredo VC et al (2014) Dose-dependent increases in p70S6K phosphorylation and intramuscular branched-chain amino acids in older men following resistance exercise and protein intake. *Physiol Rep* 2(8):e12112
- da Silva AT, de Oliveira DY, Ferreira SJ et al (2014) Sarcopenia according to the european working group on sarcopenia in older people (EWGSOP) versus Dynapenia as a risk factor for disability in the elderly. *J Nutr Health Aging* 18(5):547–553
- Dai ZL, Li XL, Xi PB et al (2012) Metabolism of select amino acids in bacteria from the pig small intestine. *Amino Acids* 42:1597–1608
- Daly RM, Connell SL, Mundell NL et al (2014) Protein-enriched diet, with the use of lean red meat, combined with progressive resistance training enhances lean tissue mass and muscle strength and reduces circulating IL-6 concentrations in elderly women: a cluster randomized controlled trial. *Am J Clin Nutr* 99:899–910
- Damas F, Phillips S, Vechin FC et al (2015) A review of resistance training-induced changes in skeletal muscle protein synthesis and their contribution to hypertrophy. *Sports Med* 45(6):801–807
- Dardevet D, Rémond D, Peyron MA et al (2012) Muscle wasting and resistance of muscle anabolism: the “anabolic threshold concept” for adapted nutritional strategies during sarcopenia. *Sci World J* 2012:269531
- Davila AM, Blachier F, Gotteland M et al (2013) Intestinal luminal nitrogen metabolism: role of the gut microbiota and consequences for the host. *Pharmacol Res* 68:95–107
- Deer RR, Volpi E (2015) Protein intake and muscle function in older adults. *Curr Opin Clin Nutr Metab Care* 18(3):248–253
- DeFronzo R, Bonadonna R, Ferrannini E (1992) Pathogenesis of NIDDM. A balanced overview. *Diabetes Care* 15(3):318–368
- Deutz N, Wolfe R (2013) Is there a maximal anabolic response to protein intake with a meal? *Clin Nutr* 32(2):309–313
- Devries MC, Phillips SM (2015) Supplemental protein in support of muscle mass and health: advantage whey. *J Food Sci* 80:A8–A15
- Devries MC, Breen L, Von Allmen M et al (2015) Low-load resistance training during step-reduction attenuates declines in muscle mass and strength and enhances anabolic sensitivity in older men. *Physiol Rep* 3(8):e12493
- Dickinson JM, Rasmussen BB (2011) Essential amino acid sensing, signaling, and transport in the regulation of human muscle protein metabolism. *Curr Opin Clin Nutr Metab Care* 14(1):83–88
- Dickinson JM, Fry CS, Drummond MJ et al (2011) Mammalian target of rapamycin complex 1 activation is required for the stimulation of human skeletal muscle protein synthesis by essential amino acids. *J Nutr* 141(5):856–862
- Dickinson JM, Drummond MJ, Coben JR et al (2013) Aging differentially affects human skeletal muscle amino acid transporter expression when essential amino acids are ingested after exercise 32(2):273–280
- Duan Y, Li F, Tan K, Liu H et al (2015) Key mediators of intracellular amino acids signaling to mTORC1 activation. *Amino Acids* 47(5):857–867
- Durham W, Casperson SL, Dillon EL et al (2010) Age-related anabolic resistance after endurance-type exercise in healthy humans. *FASEB J* 24:4117–4127
- Farnfield MM, Breen L, Carey KA et al (2012) Activation of mTOR signalling in young and old human skeletal muscle in response to combined resistance exercise and whey protein ingestion. *Appl Physiol Nutr Metab* 37:21–30
- Farsijani S, Morais JA, Payette H et al (2016) Relation between mealtime distribution of protein intake and lean mass loss in free-living older adults of the NuAge study. *Am J Clin Nutr* 104(3):694–703
- Fernandes T, Soci UPR, Alves CR et al (2008) Determinantes moleculares da hipertrofia do músculo esquelético mediados pelo treinamento físico: estudo de vias de sinalização. *Revista Mackenzie de Educação Física e Esporte* 7(1):169–188
- Fielding RA, Vellas B, Evans WJ et al (2011) Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. *International Working Group on Sarcopenia. J Am Med Dir Assoc* 12:249–256
- Fouillet H, Mariotti F, Gaudichon C et al (2002) Peripheral and splanchnic metabolism of dietary nitrogen are differently affected by the protein source in humans as assessed by compartmental modeling. *J Nutr* 132:125–133
- Francaux M, Demeulder B, Naslain D et al (2016) Aging reduces the activation of the mTORC1 pathway after resistance exercise and protein intake in human skeletal muscle: potential role of REDD1 and impaired anabolic sensitivity. *Nutrients* 8:47
- Fry CS, Drummond MJ, Glynn E et al (2011) Aging impairs contraction-induced human skeletal muscle mTORC1 signaling and protein synthesis. *Skelet Muscle* 1(1):11
- Fujita S, Rasmussen BB, Cadenas JG et al (2006) Effect of insulin on human skeletal muscle protein synthesis is modulated by insulin-induced changes in muscle blood flow and amino acid availability. *Am J Physiol Endocrinol Metab* 291:E745–E754
- Fujita S, Glynn EL, Timmerman KL et al (2009) Supraphysiological hyperinsulinemia is necessary to stimulate skeletal muscle protein anabolism in older adults: evidence of a true age-related insulin resistance of muscle protein metabolism. *Diabetologia* 52(9):1889–1898
- Fulgoni VL III (2008) Current protein intake in America: analysis of the National Health and Nutrition Examination Survey, 2003–2004. *Am J Clin Nutr* 87:1554S–1557S
- Geirsdottir O, Arnarson A, Ramel A et al (2013) Dietary protein intake is associated with lean body mass in community-dwelling older adults. *Nutr Res* 33(8):608–612
- Gibson JA, Sladen GE, Dawson AM (1976) Protein absorption and ammonia production: the effects of dietary protein and removal of the colon. *Br J Nutr* 35:61–65

- Giezenaar C, Trahair LG, Rigda R et al (2015) Lesser suppression of energy intake by orally ingested whey protein in healthy older men compared with young controls. *Am J Physiol Regul Integr Comp Physiol* 309(8):R845–R854
- Guillet C, Prod'homme M, Balage M et al (2004) Impaired anabolic response of muscle protein synthesis is associated with S6K1 dysregulation in elderly humans. *FASEB J* 18(13):1586–1587
- Gumucio JP, Mendias CL (2013) Atrogin-1, MuRF-1, and sarcopenia. *Endocrine* 43(1):12–21
- Han SS, Kim KW, Kim KI et al (2010) Lean mass index: a better predictor of mortality than body mass index in elderly Asians. *J Am Geriatr Soc* 58:312–317
- Haran PH, Rivas DA, Fielding RA (2012) Role and potential mechanisms of anabolic resistance in sarcopenia. *J Cachexia Sarcopenia Muscle* 3:157–162
- Hartman JW, Tang JE, Wilkinson SB et al (2007) Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weightlifters. *Am J Clin Nutr* 86:373–381
- Hector AJ, Marcotte GR, Churchward-Venne TA et al (2015) Whey protein supplementation preserves postprandial myofibrillar protein synthesis during short-term energy restriction in overweight and obese adults. *J Nutr* 145:246–252
- Houston D, Nicklas B, Ding J, Health ABC Study et al (2008) Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *Am J Clin Nutr* 87(1):150–155
- Ijssennagger N, Belzer C, Hooiveld GJ et al (2015) Proposed mechanism of how microbiota facilitates heme-induced compensatory hyperproliferation. *PNAS* 112:10038–10043
- Janssen I (2011) The epidemiology of sarcopenia. *Clin Geriatr Med* 27:355–363
- Johnstone AM, Murison SD, Duncan JS et al (2005) Factors influencing variation in basal metabolic rate include fat-free mass, fat mass, age, and circulating thyroxine but not sex, circulating leptin, or triiodothyronine. *Am J Clin Nutr* 82:941–948
- Jurca R, Lamonte M, Barlow C et al (2005) Association of muscular strength with incidence of metabolic syndrome in men. *Med Sci Sports Exerc* 37(11):1849–1855
- Katsanos CS, Kobayashi H, Sheffield-Moore M et al (2005) Aging is associated with diminished accretion of muscle proteins after the ingestion of a small bolus of essential amino acids. *Am J Clin Nutr* 82:1065–1073
- Katsanos CS, Kobayashi H, Sheffield-Moore M et al (2006) A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. *Am J Physiol Endocrinol Metab* 291:E381–E387
- Kim IY, Schutzler S, Schrader A et al (2015) Quantity of dietary protein intake, but not pattern of intake, affects net protein balance primarily through differences in protein synthesis in older adults. *Am J Physiol Endocrinol Metab* 308(1):E21–E28
- Kimball SR, Jefferson LS (2006) New functions for amino acids: effects on gene transcription and translation. *Am J Clin Nutr* 83(2):500S–507S
- Kumar V, Selby A, Rankin D et al (2009) Age-related differences in the dose-response relationship of muscle protein synthesis to resistance exercise in young and old men. *J Physiol* 587:211–217
- Lan A, Andriamihaja M, Blouin JM et al (2015) High-protein diet differently modifies intestinal goblet cell characteristics and mucosal cytokine expression in ileum and colon. *J Nutr Biochem* 26(1):91–98
- Lancha AH Jr, Pereira-Lancha LO (2012) *Nutrição e Metabolismo Aplicados à Atividade Motora*. Atheneu, Sao Paulo
- Landi F, Liperoti R, Russo A et al (2012) Sarcopenia as a risk factor for falls in elderly individuals: results from the il SIRENTE study. *Clin Nutr* 31(5):652–658
- Landi F, Calvani R, Tosato M et al (2016) Anorexia of aging: risk factors, consequences, and potential treatments. *Nutrients* 8(2):69
- Layman DK, Anthony TG, Rasmussen BB et al (2015) Defining meal requirements for protein to optimize metabolic roles of amino acids. *Am J Clin Nutr* 101:1330S–1338S
- Liu X, Blouin JM, Santacruz A et al (2014) High-protein diet modifies colonic microbiota and luminal environment but not colonocyte metabolism in the rat model: the increased luminal bulk connection. *Am J Physiol* 307(4):G459–G470
- Loenneke JP, Loprinzi PD, Murphy CH et al (2016) Per meal dose and frequency of protein consumption is associated with lean mass and muscle performance. *Clin Nutr*. doi:10.1016/j.clnu.2016.04.002
- Luiking Y, Deutz N, Jakel M et al (2005) Casein and soy protein meals differentially affect whole-body and splanchnic protein metabolism in healthy humans. *J Nutr* 135:1080–1087
- Malafarina V, Uriz-Otano F, Iñiesta R et al (2013) Effectiveness of nutritional supplementation on muscle mass in treatment of sarcopenia in old age: a systematic review. *J Am Med Dir Assoc* 14(1):10–17
- Mamerow MM, Mettler JA, English KL et al (2014) Dietary protein distribution positively influences 24-h muscle protein synthesis in healthy adults. *J Nutr* 144:876–880
- Marckmann P, Osther P, Pedersen A et al (2015) High-protein diets and renal health. *J Renal Nutr* 25:1–5
- Markofskia MM, Dickinson JM, Drummond MJ et al (2015) Effect of age on basal muscle protein synthesis and mTORC1 signaling in a large cohort of young and older men and women. *Exp Gerontol* 65:1–7
- McGlory C, Phillips SM (2015) Exercise and the regulation of skeletal muscle hypertrophy. *Prog Mol Biol Transl Sci* 135:153–173
- Meisel H, Bockelmann W (1999) Bioactive peptides encrypted in milk proteins: proteolytic activation and trophic-functional properties. *Antonie Van Leeuwenhoek* 76:207–215
- Mitchell CJ, Churchward-Venne TA, West DW et al (2012) Resistance exercise load does not determine training-mediated hypertrophic gains in young men. *J Appl Physiol* 113:71–77
- Mitchell CJ, Churchward-Venne TA, Bellamy L et al (2013) Muscular and systemic correlates of resistance training-induced muscle hypertrophy. *PLoS One* 8:e78636
- Mitchell CJ, Churchward-Venne TA, Parise G (2014) Acute post-exercise myofibrillar protein synthesis is not correlated with resistance training-induced muscle hypertrophy in young men. *PLoS One* 9(2):e89431
- Mitchell CJ, Della Gatta PA, Petersen AC et al (2015a) Soy protein ingestion results in less prolonged p70S6 kinase phosphorylation compared to whey protein after resistance exercise in older men. *J Int Soc Sports Nutr* 12:6
- Mitchell WK, Phillips BE, Williams JP et al (2015b) A dose—rather than delivery profile-dependent mechanism regulates the “musclefull” effect in response to oral essential amino acid intake in young men. *J Nutr* 145(2):207–214
- Mitchell WK, Phillips BE, Williams JP et al (2015c) The impact of delivery profile of essential amino acids upon skeletal muscle protein synthesis in older men. Clinical efficacy of pulse vs. bolus supply. *Am J Physiol Endocrinol Metab* 309:E450–E457
- Moore DR, Robinson MJ, Fry JL et al (2009) Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr* 89:161–168
- Moore D, Churchward-Venne T, Witard O et al. (2015) Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *J Gerontol A Biol Sci Med Sci* 70(1):57–62
- Mouillé B, Robert V, Blachier F (2004) Adaptive increase of ornithine production and decrease of ammonia metabolism in rat colonocytes after hyperproteic diet ingestion. *Am J Physiol*

- Gastrointest Liver Physiol 287(2):G344–G351. doi:10.1152/ajpgi.00445.2003
- Mu C, Yang Y, Luo Z et al (2015) Metabolomic analysis reveals distinct profiles in the plasma and urine of rats fed a high-protein diet. *Amino Acids* 47:1225–1238
- Murphy CH, Churchward-Venne TA, Mitchell CJ et al (2015) Hypoenergetic diet-induced reductions in myofibrillar protein synthesis are restored with resistance training and balanced daily protein ingestion in older men. *Am J Physiol Endocrinol Metab* 308(9):E734–E743
- Murphy CH, Oikawa SY, Phillips SM et al (2016) Dietary protein to maintain muscle mass in aging: a case for per-meal protein recommendations. *J Frailty Aging* 5:49–58
- Nair KS (2005) Aging muscle. *Am J Clin Nutr* 81:953–963
- Norton LE, Layman DK, Bunpo P et al (2009) The leucine content of a complete meal directs peak activation but not duration of skeletal muscle protein synthesis and mammalian target of rapamycin signaling in rats. *J Nutr* 139:1103–1109
- Norton C, Toomey C, McCormack WG et al (2016) Protein supplementation at breakfast and lunch for 24 weeks beyond habitual intakes increases whole-body lean tissue mass in healthy older adults. *J Nutr* 146:65–69
- Paddon-Jones D, Rasmussen B (2009) Dietary protein recommendations and the prevention of sarcopenia. *Curr Opin Clin Nutr Metab Care* 12:86–90
- Paddon-Jones D, Sheffield-Moore M, Katsanos CS et al (2006) Differential stimulation of muscle protein synthesis in elderly humans following isocaloric ingestion of amino acids or whey protein. *Exp Gerontol* 41:215–219
- Pasiakos SM, Margolis LM, Orr JS (2015) Optimized dietary strategies to protect skeletal muscle mass during periods of unavoidable energy deficit. *FASEB J* 29(4):1136–1142
- Pennings B, Boirie Y, Senden JM et al (2011a) Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men. *Am J Clin Nutr* 93:997–1005
- Pennings B, Koopman R, Beelen M et al (2011b) Exercising before protein intake allows for greater use of dietary protein-derived amino acids for de novo muscle protein synthesis in both young and elderly men. *Am J Clin Nutr* 93(2):322–331
- Pennings B, Groen B, de Lange A et al (2012) Amino acid absorption and subsequent muscle protein accretion following graded intakes of whey protein in elderly men. *Am J Physiol Endocrinol Metab* 302(8):E992–E999
- Phillips SM (2004) Protein requirements and supplementation in strength sports. *Nutrition* 20:689–695
- Phillips SM (2009) Physiologic and molecular bases of muscle hypertrophy and atrophy: impact of resistance exercise on human skeletal muscle (protein and exercise dose effects). *Appl Physiol Nutr Metab* 34:403–410
- Phillips SM (2014) A brief review of critical processes in muscular hypertrophy. *Sports Med* 44(1):S71–S77
- Phillips SM, Tang JE, Moore DR (2009) The role of milk- and soy based protein in support of muscle protein synthesis and muscle protein accretion in young and elderly persons. *J Am Coll Nutr* 28(4):343–354
- Poortmans JR, Dellalieux O (2000) Do regular high protein diets have potential health risks on kidney function in athletes? *Int J Sport Nutr Exerc Metab* 10:28–38
- Poortmans JR, Carpentier A, Pereira-Lancha LO et al (2012) Protein turnover, amino acid requirements and recommendations for athletes and active populations. *Braz J Med Biol Res* 45(10):875–890
- Ramel A, Arnarson A, Geirsdottir OG et al (2013) Glomerular filtration rate after a 12-wk resistance exercise program with post-exercise protein ingestion in community dwelling elderly. *J Nutrition* 29(5):719–723
- Rasmussen BB, Fujita S, Wolfe RR et al (2006) Insulin resistance of muscle protein metabolism in aging. *FASEB J* 20:768–769
- Reitelseder S, Agergaard J, Doessing S et al (2011) Whey and casein labeled with L-[1-13C]leucine and muscle protein synthesis: effect of resistance exercise and protein ingestion. *Am J Physiol Endocrinol Metab* 300(1):E231–E242
- Rennie MJ (2009) Anabolic resistance: the effects of aging, sexual dimorphism, and immobilization on human muscle protein turnover. *Appl Physiol Nutr Metab* 34:377–381
- Rennie MJ, Wackerhage H, Spangenburg EE et al (2004) Control of the size of the human muscle mass. *Annu Rev Physiol* 66:799–828
- Riddle ES, Stipanuk MH, Thalacker-Mercer AE (2016) Amino acids in healthy aging. *Front Biosci* 1:326–350
- Rieu I, Balage M, Sornet C et al (2006) Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia. *J Physiol* 575:305–315
- Robinson MJ, Burd NA, Breen L et al (2013) Dose-dependent responses of myofibrillar protein synthesis with beef ingestion are enhanced with resistance exercise in middle-aged men. *Appl Physiol Nutr Metab* 38:120–125
- Rondanelli M, Klersy C et al (2016) Whey protein, amino acids, and vitamin D supplementation with physical activity increases fat-free mass and strength, functionality, and quality of life and decreases inflammation in sarcopenic elderly. *Am J Clin Nutr* 103(3):830–840
- Rosenberg IH (1997) Sarcopenia: origins and clinical relevance. *J Nutr* 127:990S–991S
- Ruiz J, Sui X, Lobelo F et al (2008) Association between muscular strength and mortality in men: prospective cohort study. *BMJ* 337:a439
- Russell WR et al (2011) High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health. *Am J Clin Nutr* 93:1062–1072
- Saxton RA, Knockenbauer KE, Wolfson RL et al (2016) Structural basis for leucine sensing by the Sestrin2-mTORC1 pathway. *Science* 351(6268):53–58
- Scott D, Hayes A, Sanders K et al (2014) Operational definitions of sarcopenia and their associations with 5-year changes in falls risk in community-dwelling middle-aged and older adults. *Osteoporosis Int* 25(1):187–193
- Smith K, Reynolds N, Downie S et al (1998) Effects of flooding amino acids on incorporation of labeled amino acids into human muscle protein. *Am J Physiol* 275(1 Pt 1):E73–E78
- Tang JE, Phillips SM (2009) Maximizing muscle protein anabolism: the role of protein quality. *Curr Opin Clin Nutr Metab Care* 12:66–71
- Terzis G, Georgiadis G, Stratakis G et al (2008) Resistance exercise-induced increase in muscle mass correlates with p70S6 kinase phosphorylation in human subjects. *Eur J Appl Physiol* 102:145–152
- Ticinesi A, Meschi T, Lauretani F et al (2016) Nutrition and inflammation in older individuals: focus on vitamin D, n-3 polyunsaturated fatty acids and whey proteins. *Nutrients* 8(4):186
- Tieland M, Dirks M, van der Zwaluw N et al (2012a) Protein supplementation increases muscle mass gain during prolonged resistance-type exercise training in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 13(8):713–719
- Tieland M, van de Rest O, Dirks M et al (2012b) Protein supplementation improves physical performance in frail elderly people: a randomized, double-blind, placebo-controlled trial. *J Am Med Dir Assoc* 13:720–726

- Timmerman KL, Dhanani S, Glynn EL et al (2012) A moderate acute increase in physical activity enhances nutritive flow and the muscle protein anabolic response to mixed nutrient intake in older adults. *Am J Clin Nutr* 95:1403–1412
- Tipton KD, Gurki BE, Matin S et al (1999) Nonessential amino acids are not necessary to stimulate net muscle protein synthesis in healthy volunteers. *J Nutr Biochem* 10:89–95
- Velazquez AMC, Irigoyen CM, Delgadillo VJ et al (2013) The relationship between sarcopenia, undernutrition, physical mobility and basic activities of daily living in a group of elderly women of Mexico City. *Nutr Hosp* 28(2):514–521
- Volek JS et al (2013) Whey protein supplementation during resistance training augments lean body mass. *J Am Coll Nutr* 32(2):122–135
- Volpi E, Mittendorfer B, Wolf SE et al (1999) Oral amino acids stimulate muscle protein anabolism in the elderly despite higher first-pass splanchnic extraction. *Am J Physiol* 277(3:1):513–520
- Volpi E, Kobayashi H, Sheffield-Moore M et al (2003) Essential amino acids are primarily responsible for the amino acid stimulation of muscle protein anabolism in healthy elderly adults. *Am J Clin Nutr* 78:250–258
- Volpi E, Campbell WW, Dwyer JT et al (2013) Is the optimal level of protein intake for older adults greater than the recommended dietary allowance? *J Gerontol A Biol Sci Med Sci* 68:677–681
- Walker DK, Dickinson JM, Timmerman KL et al (2011) Exercise, amino acids, and aging in the control of human muscle protein synthesis. *Med Sci Sports Exerc* 43:2249–2258
- Wall B, van Loon L (2013) Nutritional strategies to attenuate muscle disuse atrophy. *Nutr Rev* 71(4):195–208
- Wall BT, Gorissen SH, Pennings B et al (2015) Aging is accompanied by a blunted muscle protein synthetic response to protein ingestion. *PLoS One* 10(11):e0140903
- Walrand S, Gryson C, Salles J et al (2016) Fast-digestive protein supplement for ten days overcomes muscle anabolic resistance in healthy elderly men. *Clin Nutr* 35:660–668
- Welle S, Thornton C, Jozefowicz R et al (1993) Myofibrillar protein synthesis in young and old men. *Am J Physiol* 264:E693–E698
- West D, Burd N, Coffey V et al (2011) Rapid aminoacidemia enhances myofibrillar protein synthesis and anabolic intramuscular signaling responses after resistance exercise. *Am J Clin Nutr* 94(3):795–803
- Wilkes EA, Selby AL, Atherton PJ et al (2009) Blunting of insulin inhibition of proteolysis in legs of older subjects may contribute to age-related sarcopenia. *Am J Clin Nutr* 90:1343–1350
- Wilkinson SB, Tarnopolsky MA, Macdonald MJ et al (2007) Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am J Clin Nutr* 85:1031–1040
- Witard OC, Turner JE, Jackman SR et al (2014) High dietary protein restores overreaching induced impairments in leukocyte trafficking and reduces the incidence of upper respiratory tract infection in elite cyclists. *Brain Behav Immun* 39:211–319
- Wolfe RR (2006) The underappreciated role of muscle in health and disease. *Am J Clin Nutr* 84:475–478
- Wolfe RR (2012) The role of dietary protein in optimizing muscle mass, function and health outcomes in older individuals. *Br J Nutr* 108(2):S88–S93
- Wolfe RR, Miller SL (2008) The recommended dietary allowance of protein: a misunderstood concept. *JAMA* 299:2891–2893
- Wolfe RR, Miller S, Miller K (2008) Optimal protein intake in the elderly. *Clin Nutr* 27(5):675–684
- Wolfson RL, Chantranupong L, Saxton RA et al (2016) Sestrin2 is a leucine sensor for the mTORC1 pathway. *Amino Acids* 351(6268):43–48
- Wu G (2016) Dietary protein intake and human health. *Food Funct* 7:1251–1265
- Yang Y, Breen L, Burd N et al (2012a) Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr* 108(10):1780–1788
- Yang Y, Churchward-Venne TA, Burd NA et al (2012b) Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men. *Nutr Metab (Lond)* 9(1):57
- Yarasheski KE, Zachwieja JJ, Bier DM (1993) Acute effects of resistance exercise on muscle protein synthesis rate in young and elderly men and women. *Am J Physiol* 265:E210–E214